

Is Chemical Equilibrium achieved in Collisions of Small Systems at the SPS?

Ingrid Kraus* and Helmut Oeschler

Institut für Kernphysik, Technische Universität Darmstadt, Germany

E-mail: Ingrid.Kraus@cern.ch, H.Oeschler@gsi.de

Krzysztof Redlich

Institute of Theoretical Physics, University of Wrocław, Poland

E-mail: Redlich@ift.uni.wroc.pl

Ultrarelativistic nucleus-nucleus collisions are investigated with the goal to study the properties of strongly interacting matter under extreme conditions of high energy density. Hadron multiplicities can provide information on the nature of the medium from which they are originating. The Statistical Model was recognised as a powerful approach to describe particle yields established at chemical decoupling. In general, this model assumes that at freeze-out the collision fireball appears as a statistical system in thermal and chemical equilibrium. The ensembles are constrained by charge conservation laws.

The Statistical Model has to be formulated in the canonical ensemble with respect to strangeness conservation if the number of strange particles becomes small. However, the canonical suppression under the assumption of strangeness chemical equilibrium in the whole fireball volume was found to be not sufficient to reproduce observed yields. Two approaches have been proposed to modify the model. First, a non-equilibrium factor γ_s was introduced in canonical and grand-canonical ensembles as the additional fit parameter to account for the suppressed strange particle phase-space. Here we focus on the second method: The model is extended by correlation volumes which restrict the strangeness chemical equilibrium only to certain subvolumes of the system.

In this work we report on the analysis of experimental data on particle production from p+p and central C+C, Si+Si and Pb+Pb collisions at the top SPS energy within the Statistical Model. The abundances of strange particles, in particular in the small systems, are found to be below the expectation of the Statistical Model formulated in the canonical ensemble. Therefore, we introduce strangeness equilibrated subvolumes. The canonical strangeness suppression in these correlated clusters accounts successfully for the smaller production of strange particles. The system size dependence of the correlation volume and of the thermal parameters are presented.

International Europhysics Conference on High Energy Physics

July 21st - 27th 2005

Lisboa, Portugal

*Speaker.




Figure 1: Left: Integrated π^- yield and particle ratios from Pb+Pb collisions at $\sqrt{s_{NN}} = 17.3$ AGeV together with model fits (a) (crosses), (b) (stars) and (c) (squares). The lower panel of the left side shows the deviation of the model fits to data. Middle and right: Midrapidity (squares) and 4π (circles) $\bar{\Lambda}/\Lambda$ ratio (middle) and K^-/π^- ratio (right) in p+p and central C+C, Si+Si and Pb+Pb collisions, normalised to the Pb+Pb measurement.

1. Introduction

Considering particle production in heavy ion collisions, a particular role has been attributed to strange particles because strangeness was predicted to be a sensitive probe of the properties of QCD matter. The Statistical Model [1] is very successful in describing the chemical composition of the final state of collisions over a wide range of incident energies. However, without an additional strangeness undersaturation factor, γ_s , Hadron Gas Models hardly reproduce the data from small colliding systems nor from reactions at the smaller collision energies. Here we investigate the influence of an alternative assumption, described in Section 2, on the model predictions.

In this work we investigate the system size dependence of strangeness production within Statistical Model. Experimental results at the top SPS energy, $\sqrt{s_{NN}} = 17.3$ AGeV, from p+p and central C+C, Si+Si and Pb+Pb collisions are studied [2]. Since only a limited number of hadrons were analysed in the small systems, we restricted our study on a consistent set of data. The investigated yield and ratios are demonstrated in Fig. 1 on the left hand side. Both, midrapidity densities and integrated yields were considered. Only the missing midrapidity data from p+p interactions make an exception.

2. The model description and the basic assumptions

The Statistical Model is applicable if the particles observed in high energy collisions are originating from a thermal fireball. This fireball of volume V is characterised by thermal parameters, the temperature T and baryon chemical potential μ_B , which are the same all over the system. Furthermore, we assume that strangeness is strongly correlated and appears in chemical equilibrium only within subvolumes V_C of the system. Consequently there are two volume parameters in the model. The overall volume of the system, which determines the particle yields at fixed density, and the correlation (cluster) volume V_C , which enters through the canonical suppression factor and reduces the densities of strange particles. Assuming spherical geometry the volume V_C is parameterised by




Figure 2: Chemical freeze out temperature T (left) and baryon chemical potential μ_B (right) from fits to midrapidity densities (circles) and integrated yields (squares) from p+p and central C+C, Si+Si and Pb+Pb collisions, derived with model (c). Please note the suppressed zero in the left panel.

the radius R_C which serves as a free parameter and defines the range of local equilibrium. A particle with strangeness quantum number N_S can appear anywhere in the volume V , however it has to be accompanied within the subvolume V_C by other particles carrying strangeness $-N_S$ to conserve strangeness exactly.

Three different model settings are compared: (a) an equilibrium ansatz, which uses only the temperature T and the baryon chemical potential μ_B to describe the data, (b) additionally the strangeness undersaturation factor γ_S is introduced in the standard way (multiplicative factor for each (anti)strange valence quark), (c) subvolumes with correlated strangeness production are generated by the cluster radius R_C . All Statistical Model results presented in these proceedings were achieved with the THERMUS package [3].

3. Results

The results of the three model settings together with Pb+Pb data are shown on the left hand side of Fig. 1. While the option (a) with completely equilibrated strangeness abundance fails (in particular for the small systems, not shown here), both, the approach with γ_S and the one with R_C yield comparable good descriptions of the data. These two models result in freeze-out temperatures T which are barely dependent on the system size (see left hand side of Fig. 2 for model (c)). The baryon chemical potential is flat as a function of system size if total yields are considered. In contrast, μ_B is decreasing towards smaller systems in fits to midrapidity densities (Fig. 2 left hand side). This directly reflects the $\bar{\Lambda}/\Lambda$ ratio which indicates stronger stopping in larger systems (Fig. 1, middle).

The cluster radius, R_C , varies between 0.7 fm and 1.4 fm in all data sets under study (left hand side of Fig. 3). Larger radii in the fits to midrapidity data are correlated with the experimental observation of larger K/π ratios compared to 4π yields (Fig. 1, right hand side). The increase of strange to non-strange particle ratios by a factor of 2 from p+p to Pb+Pb reactions is reflected in a variation of the subvolume radius from about 0.7 to 1.4 fm.

In the data set considered here, the fireball radius R at freeze-out is determined by the multiplicity of π^- mesons. The smaller midrapidity densities cause smaller radii R than the integrated yields. In the comparison of the cluster to the fireball radius it becomes clear that R_C has a significantly smaller system size dependence than R (right hand side of Fig. 3). In the smallest system, this




Figure 3: Cluster radius R_C as a function of the number of participants (left) and as a function of the fireball radius R (right) from fits to midrapidity densities (circles) and integrated yields (squares) from p+p and central C+C, Si+Si and Pb+Pb collisions. The line in the right panel indicates $R_C = R$.

correlation length is almost as large as the fireball radius. In contrast, with increasing system size, the subvolumes with locally conserved strangeness are up to six times smaller than the entire fireballs. In this respect, the fact that R_C is of the order of 1 fm and only weakly increasing with system size is an astonishing result and requires further studies. From this analysis we conclude that the transition from suppressed to saturated strangeness production happens in a very narrow range of the correlation length R_C . This behavior is also known from percolation model calculations [4].

To summarise, the arbitrary undersaturation factor γ_s in the Statistical Model can be replaced by the concept of strangeness equilibrated subvolumes which through canonical suppression reproduces successfully the experimental data.

References

- [1] P. Braun-Munzinger, K. Redlich and J. Stachel, *Particle Production in Heavy Ion Collisions in Quark Gluon Plasma 3*, eds. R. C. Hwa and X.-N. Wang, World Scientific Publishing, [nucl-th/0304013].
- [2] I. Kraus, *System size dependence of strange particle yields and spectra at $\sqrt{s_{NN}} = 17.3$ GeV*, *J. Phys. G* **31** (2005) S147 [nucl-ex/0412050]. A. M. Rossi et al., *Experimental Study of the Energy Dependence in Proton-Proton inclusive Reactions*, *Nucl. Phys. B* **84** (1975) 269. S. V. Afanasiev et al., *Production of ϕ -mesons in p+p, p+Pb and central Pb+Pb collisions at $E_{\text{beam}} = 158$ A GeV*, *Phys. Lett. B* **491** (2000) 59. T. Anticic et al., *Λ and $\bar{\Lambda}$ Production in Central Pb-Pb Collisions at 40, 80 and 158A GeV*, *Phys. Rev. Lett.* **93** (2004) 022302 [nucl-ex/0311024]. C. Alt et al., *System-Size Dependence of Strangeness Production in Nucleus-Nucleus Collisions at $\sqrt{s_{NN}} = 17.3$ GeV*, *Phys. Rev. Lett.* **94** (2005) 052301 [nucl-ex/0406031]. S. V. Afanasiev et al., *Energy dependence of pion and kaon production in central Pb+Pb collisions*, *Phys. Rev. C* **66** (2002) 054902.
- [3] S. Wheaton and J. Cleymans, *THERMUS - A Thermal Model Package for ROOT*, hep-ph/0407174.
- [4] M. Nardi and H. Satz, *String clustering and J/ψ suppression in nuclear collisions*, *Phys. Lett. B* **442** (1998) 14.